

Comment on “Estimating $1/f^\alpha$ scaling exponent from short time series”

P. F. Góra

*M. Smoluchowski Institute of Physics, Jagellonian University, Reymonta 4,
30-059 Kraków, Poland*

Abstract

A recently proposed method (O. Miramontes, P. Rohani, *Physica D* 166 (2002) 147) for estimating the scaling exponent in very short time series may give wrong results, especially in case of undersampled data.

Key words: Time-series, Noise parameter estimation

PACS: 05.45.Tp, 87.10.+e

Estimating the scaling exponent in the power spectrum of a signal is one of the most important tasks in the analysis of real-world temporal sequences. The point of time series analysis is not making a statement about the signal in question, but getting insight of the (unknown) mechanism responsible for generating the signal. One may say that the time series is usually not very interesting, but its generating mechanism is. The scaling exponent is one of the quantities that are calculated in order to get a better understanding of this mechanism. Many techniques have been developed for this job; see Ref. [1] for a recent review. It is particularly difficult to estimate the scaling exponent if the time series is very short, which is frequent for biological, medical and ecological data. Indeed, it is not clear whether the notion of the scaling exponent has any well-defined meaning in this case.

Short time series are usually characterized by a large sampling time, T . This is easy to understand: The series is short because it takes long time to gather the data, and it takes long time to gather the data because the sampling time is large and cannot be easily shortened. In a time series only the frequencies up to the Nyquist frequency $f_N = 1/(2T)$ are present [2]. Any possible power scaling that is discovered within the series is necessarily restricted to the Nyquist interval $[0, f_N]$. This is all the data can tell — there are no scientific grounds

Email address: gora@if.uj.edu.pl (P. F. Góra).

to claim that the scaling extends beyond the Nyquist frequency. Because T is large, f_N is small. Therefore, attributing the power scaling, if one is detected in such a series, to noise is misleading at best, as in a true noise the power scaling extends to many (theoretically to infinitely many) orders of magnitude.

When dealing with a short time series, one is also faced with another, even more acute problem: the Nyquist interval is sparsely covered. There is little one can tell about the power density at frequencies that are not multiplies of $1/(2NT)$, where N is the length of the series, and it is not clear whether the very notion of a power scaling can be used in such situations. Any attempts to interpolate between frequencies directly accessible from the data merely scale down the behaviour observed for these frequencies to those that are not directly accessible. This approach may work only if one knows *a priori* that the process used to generate the data is governed by a power law.

A method published recently in Ref. [3] suffers from all these deficiencies. The method, named “multiple segmenting method” (MSM), consists of two steps: First, one calculates scaling exponents for each segment of length 2^s , $s \geq 3$, and calculates the average scaling exponents for each s ; denote these averages by $g(n)$ with $n = 2^s$. In this step pseudo-replicates of the original data are created. These pseudo-replicates have the same short-time, or large-frequency, correlations as the original series, and the process of replicating increases their statistical significance. However, any accidental correlations among the data are enhanced as well. Second, the relation

$$g(n) = a + \frac{b}{\sqrt{n}} \quad (1)$$

is fitted to the averages calculated in the first step, and the scaling exponent for the whole series is calculated from (1) by using results of the fit and $n = N$, where N is the actual length of the time series. The equation (1) has not been derived in any rigorous way, only guessed from the example data.

The authors of Ref. [3] claim that MSM gives correct results for time series as short as $N = 47$ terms, but offer little proof to this claim. They present a number of examples with artificially created time series of moderate length $N = 400$, almost an order of magnitude larger than 47. The time series used had, by construction, built-in power scaling, and so did all their subseries. In these cases the MSM method discovered the scaling exponents that we beforehand knew were present in the data. Two out of four real-world examples presented had time series even longer ($N = 2000$ and $N = 1024$, respectively), where the power-type behaviour could be detected and the corresponding scaling exponents determined by conventional methods. There were only two really short ($N = 47$) time series of unknown properties discussed. The MSM method, when applied to these two series, gave some numbers as the scaling




Fig. 1. A quasiperiodic function (2) (thin line) and a time series resulting from undersampling the function (boxes).




Fig. 2. MSM applied to the data from Fig. 1. The thin line connects the averages for $n = 8, 16, 32$.

n	$g_1(n)$	$g_2(n)$	$g_3(n)$
8	0.66	-0.26	-1.25
16	-0.50	-0.33	-1.52
32	-0.59	-0.16	-1.05
47	-0.98	—	—

Table 1

Results of MSM method applied to the three example time series: $g_1(n)$ — the undersampled quasiperiodic function from Fig. 1, $g_2(n)$ — the undersampled AR(1) process from Fig. 3, $g_3(n)$ — first 47 terms of the full AR(1) series. The last row, second column, shows the result of applying the formula (1).

exponents, but it is not clear whether these numbers are at all meaningful.

In what follows we present examples of short time series where the MSM method, when blindfoldedly applied, gives clearly misleading results.




Fig. 3. A realization of the AR(1) process (3) (thin line) and an undersampled substring of this signal (boxes). The undersampled series is our second example time series, and the first 47 terms of the series used to draw the thin line is the third one.

As a first example, we take a quasiperiodic function

$$x(t) = \sum_{i=1}^3 a_i \sin(\omega_i t), \quad (2)$$

with $a_1 = 0.8$, $\omega_1 = 5$, $a_2 = 0.6$, $\omega_2 = 20\sqrt{3}$, $a_3 = -0.8$, $\omega_3 = 30\sqrt{3}$, sampled with a time step $T = 7.9/46 \simeq 0.17174$ to give $N = 47$. This time series is clearly undersampled, and many of the high-frequency features of the function (2) are lost, cf. Fig 1. When we apply the MSM method to this time series, we obtain a scaling exponent of about -0.98 , cf. Fig. 2 and Table 1. This result, if taken at its face value, indicates that the time series is governed by a flicker (or pink) noise, which is obviously untrue: the microscopic mechanism responsible for this time series is a regular, quasiperiodic function, albeit undersampled.

We used a realization of the Markovian AR(1) process [4]

$$X_{n+1} = \beta X_n + (1 - \beta)\eta_n, \quad (3)$$

with $\beta = 0.69$ like in [3], to generate two next example time series. η_n is a Gaussian white noise, generated numerically by means of an algorithm published in [5]. We generated 256 terms of the series (3), and then constructed our second example time series by taking every fifth term of the original series up to $N = 47$ ($X_5, X_{10}, \dots, X_{235}$). Finally, the first consecutive 47 terms of the original series ($X_1, X_2, \dots, X_{47}$) were used as the third example. The output of the MSM method is presented in Figs. 4 and 5 and in Table 1. Note that the function (1) does not seem to be a reasonable fit to these data, and if this fit fails, MSM does not offer any other means to systematically determine the scaling exponent for the whole series. For the second example the method in-




Fig. 4. Results of the MSM method applied to the second example time series. The thin line connects the averages for $n = 8, 16, 32$.




Fig. 5. Results of the MSM method applied to the third example time series (47 terms). The thin line connects the averages for $n = 8, 16, 32$. The inset shows the power spectrum calculated from the whole 256 terms used to generate the data. The line with the slope -1.5 is not a rigorous fit — it is meant as a guide to the eye only.

dictates the scaling exponent in the range $[-0.33, -0.16]$, i.e. it recognizes the time series as being close to the white noise, which is again not true. On the other hand, for the third example the method predicts the scaling exponent in the range $[-1.52, -1.05]$, which is close (especially the lowest value) to a rough estimate for the scaling exponent for the whole 256 term time series. Note that this time series was, by construction, characterized by a power-type behaviour. The relative success of MSM in this particular case strengthens our point that *if* a power law is present in the data, scaling it down to frequencies not observed in the data may not hurt much, but a short time series does not tell *whether* such a power law is present.

One may argue that the MSM method failed for the undersampled time series only. However, undersampling can be quite common in practice. When dealing with real-world time series generated by a mechanism whose details remain

unknown, we never know whether we choose a correct sampling time. For instance, in the bacterial population data analysed in Ref. [3] the sampling time was as long as a week, while the true dynamics of the population might have been governed by processes with a much lower characteristic time, like daily changes in sunlight or temperature. Similarly, in many medical studies levels of drugs in subjects are determined on a daily basis, while physiological processes responsible for the drug absorption and decay usually proceed much faster. There are many other situations in which data are collected not in a controlled laboratory environment, but come out of real-world processes, where adjusting the sampling time may be impossible for various “technical” reasons. In case of undersampling, the power spectrum beyond the Nyquist frequency is aliased into the Nyquist interval, distorting the available power spectrum [2]. Examples presented above show that the MSM method, when applied to such data, may give wildly wrong results.

One needs to know some important details of the underlying process in order to estimate its parameters from a very short time series. Otherwise idiosyncrasies of the data are mistakenly taken for true correlations and are enhanced in the process of pseudo-replicating the time series. This may lead to a failure of the MSM method. This method appears to work tolerably well for long time series, but in this case it is inferior to conventional techniques due to its large computational costs. In our opinion, the usefulness of the MSM method is thus limited to the rare case when one knows for sure that the process used to generate the time series is governed by a power law with an unknown exponent, but the time series available is very short and cannot be easily extended and only a rough estimate of the scaling exponent is required. In other situations the MSM method is not recommended.

I would like to thank Dr. Adam Kleczkowski from Cambridge University for his helpful comments.

References

- [1] B. Pilgrim, D. T. Kaplan, *Physica D* 114 (1998) 108.
- [2] W. H. Press, S. A. Teukolsky, W. T. Vetterling, B. P. Flannery, *Numerical Recipes in Fortran/C: The Art of Scientific Computing, Second Edition* (Cambridge University Press, Cambridge, 1992)
- [3] O. Miramontes, P. Rohani, *Physica D* 166 (2002) 147.
- [4] G. E. P. Box, G. M. Jenkins, *Time Series Analysis* (Holden-Day, San Francisco, 1976)
- [5] J. F. Fernandez and C. Criado, *Phys. Rev. E* 60 (1999) 3361.